

| Fatty Acid Methyl Esters*                         |
|---|
| C12:0 (Lauric)                                    |
| C14:0 (Myristic)                                  |
| C14:1 (Myristoleic)                               |
| C16:0 (Palmitic)                                  |
| C16:1 (Palmitoleic)                               |
| C17:0 (Heptadecanoic)                             |
| C18:0 (Stearic)                                   |
| C18:1n9c (Oleic)                                  |
| C18:2n6c (Linoleic) LA                            |
| C18:3n3 ( $\alpha$ -Linolenic)                    |
| C18:3n6 ( $\gamma$ -Linolenic)                    |
| C20:0 (Arachidic)                                 |
| C20:2 (cis-11,14-Eicosadienoic)                   |
| C20:3n6 (cis-8,11,14-Eicosatrienoic)              |
| C20:4n6 (Arachidonic) AA                          |
| C20:5n3 (cis-5,8,11,14,17-Eicosapentaenoic) EPA   |
| C22:6n3 (cis-4,7,10,13,16,19-Docosahexaenoic) DHA |
| C22:0 (Behenic)                                   |
| C22:5 (cis-7,10,13,16,19-Docosapentaenoic) DPA    |
| C24:0 (Lignoceric)                                |
| C24:1n9 (Nervonic)                                |

\*Measurement of total fatty acids as free fatty acids (FFAs) and esterified fatty acids such as glycerolipids (mono-, di- and triglycerides) and glycerophospholipids (eg. Phosphatidylcholines, phosphatidylethanolamines, etc.)